

The Massive Wolf-Rayet Binary HDE 318016 (=WR 98)

Roberto C. Gamen¹ and Virpi S. Niemela²

Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata. Paseo del Bosque s/n, B1900FWA, La Plata, Argentina.

Abstract. We present the discovery of OB type absorption lines superimposed to the emission line spectrum, and the first double-lined orbital elements for the massive Wolf-Rayet binary HDE 318016 (=WR 98), a spectroscopic binary in a circular orbit with a period of 47.825 days. The semiamplitudes of the orbital motion of the emission lines differ from line to line, indicating mass ratios between 1 and 1.7 for $\mathcal{M}_{WR}/\mathcal{M}_{OB}$.

1. Introduction and Observations

HDE 318016 =WR 98 is one of the few stars with Wolf-Rayet spectrum showing both N and C emission lines. This star was found to be a single-lined binary with a period of 47.8 days with N and C emission lines moving in phase (Niemela 1991). Relevant parameters of WR 98 can be found in the recent VIIth Catalogue of galactic WR stars (van der Hucht 2001).

Here we present the results of a detailed radial velocity analysis of optical spectral lines of WR 98 showing it to be a double-lined binary.

We have obtained a total of 69 blue optical spectrograms of WR 98 at two different Observatories, namely CTIO in Chile (1980-84) and CASLEO (1997–2001) in Argentina. The observations at CTIO were performed with the Cassegrain IT spectrograph on the 1m Yale telescope, using photographic plates with fine grain emulsion as detector. The observations at CASLEO were obtained with the Cassegrain spectrographs with CCD detectors attached to the 2.15m telescope. (CASLEO is operated under agreement between CONICET, SeCyT, and the Universities of La Plata, Córdoba and San Juan, Argentina.)

The photographic observations were digitized with a Grant engine at La Plata. The digital data were processed and measured with IRAF routines.

2. Results

Our radial velocity analysis of the optical spectral lines of WR 98 using a long term database, confirms that this star is a binary system in a circular orbit with a period of 47.825 days. Figure 1 depicts the radial velocity variations of the

¹ Fellow of CONICET, Argentina

²Visiting Astronomer, CTIO, NOAO, operated by AURA, Inc. for NSF, USA. Member of Carrera del Investigador, CIC-BA, Argentina




Figure 1. Left: $H\gamma$ and $H\beta$ absorptions observed atop of the $He II$ emissions. Right: Radial velocity variations of emission lines, and absorption lines of $H\gamma$ (circles) and $H\beta$ (triangles) phased with $P=47.825$ days. Curves represent the orbital solutions from Table 1.

spectral lines in this period. N and C emission lines in the spectrum move in phase, indicating that they are formed in the same stellar envelope (See Fig. 1).

We have also detected absorption lines upon the emissions. These absorptions correspond to a late O-type spectrum, and they move anti-phased with respect to the emission lines (see Fig. 1). Thus WR 98 is a double-lined spectroscopic binary. Orbital elements are listed in Table 1. Radial velocity variations of absorption and emission lines indicate that the mass of the WR component is similar or higher than that of the OB component (See Table 1).

Table 1. Circular Orbital Elements of WR 98 ($P=47.825$ days).

Element	N IV em.	N V em.	C III em.	He II em.	absorptions
V_0 [km s $^{-1}$]	-41 ± 4	-15 ± 3	-47 ± 2	4 ± 2	6 ± 15
K [km s $^{-1}$]	106 ± 6	109 ± 5	72 ± 3	65 ± 3	112 ± 16
$\mathcal{M}_{WN} \sin^3 i$ [$\mathcal{M}_\odot$]	27 ± 10	28 ± 10	19 ± 8	18 ± 7	
$\mathcal{M}_{OB} \sin^3 i$ [$\mathcal{M}_\odot$]	25 ± 7	27 ± 7	12 ± 3	10 ± 3	

References

- van der Hucht, K. A. 2001, *New Astronomy Reviews*, 45, 135
 Niemela, V. S. 1991, in K. A. van der Hucht & B. Hidayat (eds.), *Wolf-Rayet Stars and Interrelations with Other Massive Stars in Galaxies*, Proc. IAU Symp. 143, 201